

## Activyl® for Cats and Kittens

### Product Description:

Activyl® for Cats and Kittens is a once-a-month topical flea treatment for cats and kittens 8 weeks of age or older and 2 lbs. There are two sizes available depending on the size of your cat. Monthly treatments are recommended for optimal control and prevention of fleas. It is available in an easy to use applicator.

### Product Bullets:

- Kills fleas within 24 hours of application; starts killing fleas within 8 hours
- Kills fleas that may cause flea allergy dermatitis
- Long-lasting protection against fleas for up to 1 month.
- Breaks the flea life cycle to prevent new fleas from developing
- Fragrance-Free, Quick-Drying
- Available in 2 sizes based on your cat's weight
- Primary ingredient: Indoxacarb

### Directions for Use:

For external use on cats and kittens only. Do not use on kittens under 8 weeks of age and weighing less than 2 lbs.

- Open one foil pouch and remove applicator.
- Step 1: Hold applicator in upright position away from your face and snap tip open by bending it and folding it back on itself.
- Step 2: Part your cat's coat and place the applicator tip against the skin at the base of her skull.
- Step 3: Squeeze applicator firmly applying entire contents directly to skin. Do not allow product to run off. Do not have contact with treated area until it is completely dry.
- It is recommended to reapply every 30 days or as needed, but DO NOT apply more than 1 applicator per month.
- When application is complete, follow the Container Handling instructions under the Storage and Disposal section below.
- Wash thoroughly with soap and water after handling

**SIDE EFFECTS:** Monitor your cat after application. Accidental ingestion by the cat may cause intermittent excess salivation. Other immediate post-treatment observations may include licking the face, shaking the head and squinting. In rare cases, application site reactions (e.g. scratching, redness, hair loss) may occur. These effects usually resolve without requiring treatment. Rare cases (fewer than 1 in 1,000) of neurological signs (e.g. incoordination, tremor, impaired balance, convulsions, dilated pupils, and impaired vision) and/or vomiting and very rare cases (fewer than 1 in 10,000) of decreased appetite, lethargy, hyperactivity, or vocalization may occur. These effects will generally resolve with supportive treatment. If these or other side effects occur consult a veterinarian immediately.

**HAZARDS TO HUMANS:** Causes substantial but temporary eye injury. Harmful if absorbed through skin or swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**HAZARDS TO DOMESTIC ANIMALS:** For external use on cats and kittens only. Do not use on kittens under 8 weeks of age OR weighing less than 2 lb. Do not use on cats (male or female) intended for breeding, pregnant or nursing. Consult a veterinarian before using this product on debilitated, aged, or medicated cats. Accidental ingestion by the cat may cause intermittent excess salivation. Other immediate post-treatment observations may include licking the face, shaking the head and squinting. These effects should disappear within several minutes following exposure without requiring treatment. Individual sensitivities may occur after using any pesticide product. If signs of sensitivity occur, consult a veterinarian immediately. Certain medications can interact with pesticides.

**PHYSICAL OR CHEMICAL HAZARDS:** Flammable. Keep away from heat and open flame.

### STORAGE AND DISPOSAL:

**PESTICIDE STORAGE:** Store in a cool, dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** If empty, do not reuse the container. Place in trash or offer for recycling if available. If partly filled, call your local solid waste agency for disposal instructions. Never place unused products down any indoor or outdoor drain.