

# Prime Pac® PRRS RR

PORCINE REPRODUCTIVE AND RESPIRATORY SYNDROME VACCINE

Reproductive and Respiratory Form, Modified Live Virus



This product has been shown to be effective for the vaccination of healthy swine 3 weeks of age or older against respiratory disease caused by Porcine Reproductive and Respiratory Syndrome (PRRS) virus and female breeding age swine against reproductive disease caused by PRRS virus.

## Advantages:

- Prime Pac PRRS RR is a safe product to include in your respiratory and reproductive disease prevention protocols.
- Licensing data demonstrates good cross-protection in the face of a heterologous PRRSV challenge.
- Keeps pigs healthy in the face of a heterologous PRRSV challenge.
- Duration of Immunity against reproductive disease is at least 20 weeks.
- Convenient 1mL dose.



Code: 187813 100mL  
100 doses - 1mL dose

Code: 188024 20mL  
20 doses - 1mL dose

*Prime Pac PRRS RR is recommended for use in healthy swine from herds with a PRRS virus-positive history according to label direction. Recommended use shall be restricted to premises having a history of the disease.*

DRIVEN BY



PREVENTION™



# Prime Pac<sup>®</sup> PRRS RR

PORCINE REPRODUCTIVE AND RESPIRATORY SYNDROME VACCINE



**ADMINISTRATION/DOSAGE:** Aseptically rehydrate Prime Pac PRRS RR with the diluent supplied. Sows and gilts: Inject a single 1 mL dose intramuscularly (IM) 8 weeks prior to breeding. Piglets: Inject a single 1 mL dose IM at 3 weeks of age or older. Duration of Immunity against reproductive disease is at least 20 weeks. Duration of Immunity against respiratory disease has not been established; consultation with a veterinarian is recommended.

## PREPARATION OF THE VACCINE:

1. Do not rehydrate the vaccine until ready for use.
2. Mix only one set of vials at a time and use entire contents promptly after rehydration.
3. Using a sterile transfer needle, transfer the entire contents of sterile diluent into the bottle of freeze dried vaccine and shake the combined vaccine and diluent to suspend evenly. The vaccine is now ready for use.

**CAUTION:** Vaccine virus can shed and be transmitted to other populations of swine in direct or indirect contact with vaccinated swine. The duration of potential vaccine virus transmission may vary. Non-vaccinated pigs in contact with modified live PRRS-vaccinated pigs may seroconvert to vaccine virus. Use of the vaccine in herds intended to remain PRRS virus seronegative is contraindicated. Introduction of vaccinated pigs into herds intended to remain PRRS virus seronegative is contraindicated.

Store at 2° to 8°C (35° to 46°F). Use aseptic technique and only sterile needles and syringes to inject the vaccine. Use entire contents when first opened. Do not mix with other products. Inactivate unused contents before disposal. Do not vaccinate within 21 days before slaughter. Not for use in pregnant sows or boars. If allergic response occurs, administer epinephrine. In case of human exposure, contact a physician.

**PRESERVATIVE:** Gentamicin.

## FOR ANIMAL USE ONLY

Intervet Inc. d/b/a Merck Animal Health  
Omaha, Nebraska 68103 USA  
VLN 165A/PCN 19T1.21  
1 800 521-5767