

From the manufacturer



safe-guard®  
(fenbendazole)  
**98.7%**  
Efficacy<sup>1</sup>



**Safe for Any Stage of Gestation**  
The active ingredient in SAFE-GUARD®, fenbendazole, offers strategic deworming solutions for your animals. SAFE-GUARD® works differently than other dewormers because of this active ingredient. It goes straight to the gut, killing the parasites where they live.

**Effective Formulation**  
Fenbendazole builds up in the fat tissue of parasites. Even cattle that ingest small amounts of SAFE-GUARD over the course of a few feedings will quickly build up to a dose that is lethal to internal parasites.

**A Legacy of Trusted Protection**  
For 40 years, producers have relied on SAFE-GUARD to protect their herds from profit-eating parasites. It was powerful then, and remains powerful now – with all formulations demonstrating efficacy over 91% as determined by ongoing testing.<sup>1</sup>

**Low-dose volume paste designed for convenient, stress-free deworming.**

Each 290 gm paste cartridge deworms 12,600 lb. of cattle

safe-guard® (fenbendazole)

**Dosing Done Right**

Product	Options for Deworming Chute Side		Total Pounds Treated per Unit
	500 lb Head Treated	1,300 lb Head Treated	
1 L Drench	86	33	43,478
1 gal Drench	329	126	164,565
10 L Drench	869	334	434,780
92 g Paste	8	3	4,000
290 g Paste	24	9	12,000

**Deworm your way with SAFE-GUARD. Available in a wide variety of effective formulations.**



**IMPORTANT SAFETY INFORMATION:**

SAFE-GUARD®: Do not use in beef calves less than 2 months old, dairy calves and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves. Additionally, the following meat withdrawal and milk discard times apply:  
SAFE-GUARD Paste: Cattle must not be slaughtered for 8 days. For dairy cattle, the milk discard time is 96 hours. SAFE-GUARD Suspension: Cattle must not be slaughtered for 8 days. For dairy cattle, the milk discard time is 48 hours.  
SAFE-GUARD® ENPROAL® Type C Medicated Block: Cattle must not be slaughtered for 11 days. For use in beef cattle only.  
SAFE-GUARD® 20% Protein Type C Medicated Block: Cattle must not be slaughtered for 16 days. For use in beef cattle only.  
Safe-Guard Type A and other medicated feed products. (pellets, cubes, free-choice mineral, or free-choice liquid): Cattle must not be slaughtered for 13 days. For dairy cattle, the milk discard time is 60 hours.

<sup>1</sup> Merck Animal Health National FECRT Database.  
Copyright © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.