

From the manufacturer



Safe for Any Stage of Gestation
The active ingredient in SAFE-GUARD®, fenbendazole, offers strategic deworming solutions for your animals. SAFE-GUARD® works differently than other dewormers because of this active ingredient. It goes straight to the gut, killing the parasites where they live.





Effective Formulation
Fenbendazole builds up in the fat tissue of parasites. Even cattle that ingest small amounts of SAFE-GUARD over the course of a few feedings will quickly build up to a dose that is lethal to internal parasites.



A Legacy of Trusted Protection
For 40 years, producers have relied on SAFE-GUARD to protect their herds from profit-eating parasites. It was powerful then, and remains powerful now – with all formulations demonstrating efficacy over 91% as determined by ongoing testing.¹

Low-dose volume suspension with easy-to-use applicator for accurate, stress-free deworming.

One liter deworms 43,478 lb. of cattle 



safe-guard® (fenbendazole)

Dosing Done Right

Product	Options for Deworming Chute Side		Total Pounds Treated per Unit
	500 lb Head Treated	1,300 lb Head Treated	
1 L Drench	86	33	43,478
1 gal Drench	329	126	164,565
10 L Drench	869	334	434,780
92 g Paste	8	3	4,000
290 g Paste	24	9	12,000

Deworm your way with SAFE-GUARD. Available in a wide variety of effective formulations.



IMPORTANT SAFETY INFORMATION:

SAFE-GUARD®: Do not use in beef calves less than 2 months old, dairy calves and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves. Additionally, the following meat withdrawal and milk discard times apply:
SAFE-GUARD Paste: Cattle must not be slaughtered for 8 days. For dairy cattle, the milk discard time is 96 hours. SAFE-GUARD Suspension: Cattle must not be slaughtered for 8 days. For dairy cattle, the milk discard time is 48 hours.
SAFE-GUARD® ENPROAL® Type C Medicated Block: Cattle must not be slaughtered for 11 days. For use in beef cattle only.
SAFE-GUARD® 20% Protein Type C Medicated Block: Cattle must not be slaughtered for 16 days. For use in beef cattle only.
Safe-Guard Type A and other medicated feed products (pellets, cubes, free-choice mineral, or free-choice liquid): Cattle must not be slaughtered for 13 days. For dairy cattle, the milk discard time is 60 hours.

¹Merck Animal Health National FECRT Database.
Copyright © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.