

SAFE-GUARD®

(fenbendazole)

20% Type A Medicated Article



For the removal and control of gastrointestinal worms: round worms, adults and larvae (*Ascaridia dissimilis*), cecal worms, adults and larvae (*Heterakis gallinarum*), an important vector of *Histomonas meleagridis* (Blackhead) in growing turkeys.

ADVANTAGES:

- Proven safety
- The only in-feed dewormer approved for use in turkeys
- Excellent efficacy against both larval and adult roundworms¹
- No pre-slaughter withdrawal required. May be used until day of slaughter.²

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.



Safe-Guard® is an anthelmintic type A medicated premix containing 20% fenbendazole for use in growing turkeys.

Net Weight: 25 lb (11.34 kg)

¹ Data on file. Merck Animal Health

² Safe-Guard SAFE-GUARD® Type "A" Medicated article (Premix) [prescribing information]; 1991

SAFE-GUARD[®]

(fenbendazole)

20% Type A Medicated Article



For Animal Use Only.

Description

An anthelmintic feed additive for the control of gastrointestinal worms in growing turkeys.

Active ingredient: Fenbendazole 200 grams per kilogram (90.7 grams per pound) Inert ingredients: roughage products or roughage products and calcium carbonate; and mineral oil or soybean oil.

Indications for Use

For the removal and control of gastrointestinal worms: round worms, adults and larvae (*Ascaridia dissimilis*), cecal worms, adults and larvae (*Heterakis gallinarum*), an important vector of *Histomonas meleagridis* (Blackhead) in growing turkeys.

Important

Must be thoroughly mixed into turkey feed before use.

Directions for Mixing

Thoroughly mix 72.5 grams (0.16 lb) of Safe-Guard (fenbendazole) 20% Type A Medicated Article into each ton of complete feed to provide 16 ppm fenbendazole (use level).

It is recommended that an intermediate mix containing 1 part Safe-Guard 20% Type A Medicated Article and not less than 9 parts appropriate feed ingredients be thoroughly mixed before incorporation into the final feed.

The working premix is then blended with the complete feed mixture. Thoroughly mix both working premix and complete feed to assure complete and uniform distribution of the Safe-Guard 20% Type A Medicated Article.

Directions for Use

Fenbendazole is added to turkey feed at a concentration of 16 ppm (14.5 g/ton). The resultant complete turkey feed containing fenbendazole is then fed as the sole ration for 6 consecutive days.

Warning

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. NOT FOR USE IN HUMANS.

This product is not hazardous when used according to directions supplied. A safety data sheet (SDS) is available upon request. This and any other consumer information can be obtained by calling Merck Animal Health Customer Service at 1-800-211-3573.

Residue Warning

There is no pre-slaughter withdrawal period as Safe-Guard 20% Type A Medicated Article can be fed to day of slaughter.

Ask your Merck Animal Health sales or technical services representative to design a custom gastrointestinal worm control program to meet your needs.