

# SAFE-GUARD® AQUASOL®

## (Fenbendazole oral suspension)

Innovative manufacturing process enables easy, effective deworming for broilers, breeders, pullets and laying hens through oral administration via drinking water.

### NEW PRODUCTION TECHNOLOGY<sup>1</sup>

- Now approved for laying hens and replacement chickens intended to become laying hens
- Proprietary, innovative wet-milling technique produces highly stable suspension in drinking water
- Smaller particle size helps prevent sedimentation and clogged pipes, filters or nipple drinkers
- Easily administered by medicators or proportioners

### APPROVED FOR MAJOR INTESTINAL PARASITES<sup>2</sup>

- *Ascaridia galli* (adult stage roundworms)
- *Heterakis gallinarum* (adult stage cecal worms)

### EFFECTIVE<sup>2</sup>

- 98.8% efficacy against *A. galli* in 4 US field studies (100% in broilers)
- 92.1% efficacy against *H. gallinarum* (long lived birds)

### CONVENIENT WATER FORMULATION

- Short, 5 day treatment
- No pre-slaughter withdrawal period when used as directed
- Shelf life of 3 years unopened, 6 months after opening — multi-use container reduces waste

### SAFE<sup>2</sup>

- Proven safe when used as directed
- No adverse effects on egg production, hatchability or progeny of breeders
- No impact on water or feed intake compared to unmedicated controls

<sup>1</sup>Data on file, Merck Animal Health

<sup>2</sup>Freedom of Information Summary, NADA 141-449



Safe-Guard® AquaSol is available in 1 liter and 1 gallon presentations.

# SAFE-GUARD<sup>®</sup> AQUASOL<sup>®</sup>

## (Fenbendazole oral suspension)

For Animal Use Only.

### Description

Safe-Guard<sup>®</sup> AquaSol is a suspension concentrate containing fenbendazole, an antiparasitic. Each mL of Safe-Guard AquaSol contains 200 mg of fenbendazole.

### Indications for Use

Safe-Guard AquaSol is indicated for the treatment and control of adult *Ascaridia galli* in broiler chickens and replacement chickens and for the treatment and control of adult *A. galli* and *Heterakis gallinarum* in breeding chickens and laying hens.

### Dosage and Administration

Safe-Guard AquaSol must be administered orally to chickens via the drinking water at a daily dose of 1.0 mg/kg BW (0.454 mg/lb) for 5 consecutive days.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

### Residue Warning

No withdrawal period is required when used according to labeling.

### Storage Information

Store at room temperature 30°C (86°F). Once opened, do not store the container above 25 °C (77°F). Do not freeze.

Use within 6 months after opening. Use the medicated water within 24 hours.

For customer service, adverse effects reporting, and/or a copy of the SDS, call 1-800-211-3573